

# Pediatric Associates of Plano

## Acetaminophen

for Infants or Children  
DOSAGE: EVERY 4-6 HOURS.  
**NOT TO EXCEED 5 DOSES IN 24 HOURS.**  
*fever reducer, pain reliever*

Dose		160 mg/5 mL Liquid or Suspension		80 mg Chewable Tablets	160 mg Chewable Tablets
Weight	Age	mL	tsp	Tablet	Tablet
For most accurate dosing, use a dosing syringe or dosing spoon					
6-11 lbs	0-3 mos	1.25 mL	1/4 tsp		
12-17 lbs	4-11 mos	2.5 mL	1/2 tsp		
18-23 lbs	12-23 mos	3.75 mL	3/4 tsp		
24-35 lbs	2-3 yrs	5 mL	1 tsp	2	
36-47 lbs	4-5 yrs	7.5 mL	1 1/2 tsp	3	
48-59 lbs	6-8 yrs	10 mL	2 tsp	4	2
60-71 lbs	9-10 yrs	12.5 mL	2 1/2 tsp	5	2 1/2
72-95 lbs	11 yrs	15 mL	3 tsp	6	3
96 lbs & over	12 yrs				4

**\*\*\*LOOK CAREFULLY AT YOUR MEDICATION'S CONCENTRATION!\*\*\***

## Ibuprofen

for Infants or Children  
DOSAGE: EVERY 6-8 HOURS.  
**NOT TO EXCEED 3 DOSES IN 24 HOURS.**  
*anti-inflammatory, pain reliever,  
 fever reducer*

Dose		50 mg/1.25 mL Liquid, Drops, or Suspension	100 mg/5 mL Liquid or Suspension	50 mg Chewable Tablets	100 mg Chewable Tablets	
Weight	Age	We do not recommend using this concentration	mL	tsp	Tablet	Tablet
For most accurate dosing, use a dosing syringe or dosing spoon						
12-17 lbs	6-11 mos		2.5 mL	1/2 tsp		
18-23 lbs	12-23 mos		5 mL	1 tsp	2	1
24-35 lbs	2-3 yrs		7.5 mL	1.5 tsp	3	1 1/2
36-47 lbs	4-5 yrs		10 mL	2 tsp	4	2
48-59 lbs	6-8 yrs			2 1/2 tsp	5	2 1/2
60-71 lbs	9-10 yrs			3 tsp	6	3
72-95 lbs	11 yrs			4 tsp	8	4
96 lbs & over	12 yrs					adult dosing

**\*\*\*LOOK CAREFULLY AT YOUR MEDICATION'S CONCENTRATION!\*\*\***